

Journal of Family Medicine & Community Health

Perspective

The Role of General Practices for Monitoring and Protecting the Environment and Health. Results and Proposals of the Italian Project Aimed at Creating an "Italian Network of Sentinel Physicians for the Environment" (Rimsa) within an International Perspective

Paolo Lauriola*, Samantha Pegoraro, Alice Serafini, Vitalia Murgia, Agostino Di Ciaula, Francesco De Tommasi, Alice Rossci, Mariagrazia Santamaria, Giacomo Toffol, Fabrizio Bianchi, Roberto Romizi, Emanuele Vinci, Behrooz Behbod, Ariana Zeka, Robert Verheij,

Giovanni Leonardi, Raymond Agius

Italian National Research Council, Institute of Clinical Physiology, Unit of Environmental epidemiology and disease registries, International Society Doctors for the Environment Italy, Italy

*Corresponding author

Paolo Lauriola, Italian National Research Council, Institute of Clinical Physiology, Unit of Environmental epidemiology and disease registries, International Society Doctors for the Environment Italy, Italy, Tel: +39-335 5756961; Email: paolo.lauriola@gmail.com

Submitted: 30 July 2018
Accepted: 22 October 2018
Published: 28 October 2018

ISSN: 2379-0547 Copyright

© 2018 Lauriola et al.

OPEN ACCESS

Keywords

- Sentinel family doctors
- Environmental-health surveillance
- Climate change
- Primary health Care
- Environmental epidemiology

Abstract

Health effects due to the environment are among the most challenging concerns faced by our future. In particular global climate changes. However, the connection between the environment and human health as part of clinician's activities is rather undeveloped. We performed a literature review, finding that many Sentinel Physician Networks have been developed, but very few have dealt with environmental health issues, and almost none have considered the opportunities of targeting Family Doctors (FDs). Beginning with the Italian context, we are committed to developing a structured system, named RIMSA – Rete Italiana Medici Sentinella per l'Ambiente – (Sentinel Physicians for the Environment) as a pilot experience. Moreover, we would like to share our experience to raise interest of this model internationally. The aim is to monitor the effects of critical environmental public health issues, help raise awareness of environmental health risks among doctors, patients, and the general public, and learn how to report sentinel diagnoses or critical situations. We believe FDs can play an important role in connecting global concerns with local actions. The project has already made significant strides; three 2-day courses have been staged, training 61 FDs from across Italy; a website, a Moodle platform and an online survey have been implemented and the Sentinel Physicians for the Environment (SPE) Manual is currently being compiled. Other actions have been planned to consolidate and develop the project; the training process should be sustained and disseminated and a standard plan for reporting an environmental health concern is underway. This will represent the foundation for setting up the overall SPE network in Italy and possibly in connection with other Sentinel Practitioners Networks in Europe and globally.

INTRODUCTION

Major and minor effects of the environment on health: how to track them?

According to the World Health Organisation (WHO), about

12.6 million deaths and disabilities were attributed to the environment in 2012. About a quarter of the global burden of diseases is linked to environmental factors (22%; 95% CI 12%-32%) [1].

In particular, the health effects due to climate change will



be the most challenging concern in coming years [2], with the WHO estimating an additional 250,000 deaths/year between 2030 and 2050 due to the negative effects of climate change [3]. These figures are an underestimate of the overall health impacts of climate change, since the estimates do not include the indirect impacts of climate change on other major determinants of health, such as migration and conflict.

In addition to the health effects usually quantified through the number of deaths, emergency admissions and hospitalizations, there are many other indicators linked to climate change that could provide figures of the true impact in terms of the day-to-day life of people and on the overall organisation of healthcare services. Possible health consequences could be secondary to a number of clinical conditions that are neither necessarily lethal nor require hospitalisation such as, allergic diseases and asthma [4], hypertension [5], fluid and electrolyte disorders [6], child and adult obesity [7], type 2 diabetes and its complications [8-10], panic attacks [11], and Parkinson's disease [12]. Moreover, different countries may have their own specific concerns such as arthropod-borne infectious diseases, as in the case of disease transmitted by *Aedes albopictus*, a permanently present vector throughout Italy [13,14].

Such diseases are generally treated by Family Doctors (FDs), and are therefore not always linked with environmental health (EH) threats. This drawback can be overcome by implementing a surveillance system aimed at systematically collecting, analysing, interpreting and reporting health and environmental data, thereby translating into public health actions.

The importance of FDs as Health System gatekeepers and sentinels of the Environment and Health

In 2003 and 2009, the World Organization of National Colleges, Academies and Academic Associations of General Practitioners/Family Physicians, (WONCA) organized two international conferences on how to promote research in Family Medicine. On those occasions, it was assessed that in primary care led healthcare, 90–95% of patients were initially cared for by FDs, with 80% of health problems being managed entirely within primary care. WONCA also stated that FDs played an influential role on individual patients and communities, both in terms of individual attitudes and collective planning and choices [15].

FDs could be a very valuable source of useful data, helping inform decisions leading to effective global climate change (GCC) adaptation measures at the national and local level. With paradigmatic examples such as vector-borne diseases (VBD), which are highly variable geographically and are known as "focal diseases" [16,17], global strategies addressing GCC should be implemented locally.

As such, FDs could play a helpful role in connecting global concerns with local actions through the information they provide and their influence on individual attitudes and policy [18]. Nevertheless, it must be emphasised that a gap still exists in the activities of physicians with regard to the relationship between health and the environment in which their patients work and live.

It is therefore necessary to guarantee:

Sound science and evidence base, both to determine the proposed processes, and to evaluate their future outcomes; Clear and effective communication strategies among all stakeholders, including FDs, communities, individuals and policy-makers; Actions aimed at raising and strengthening EH awareness among physicians; Specific, effective, and motivating training of existing and future FDs in the field of EH.

Sentinel Physicians worldwide: a literature review

Traditionally "a sentinel general practice network - or sentinel network of general practitioners - can be defined as a system that keeps a watchful eye on a sample of the (overall¹) population by supplying regular and standardized reports on the incidence and the main epidemiological characteristics of specific diseases and of procedures in primary health care" [19]. More recently, sentinel FDs provide data to a central database via electronic medical records (EMRs) to work out meaningful epidemiological information.

We reviewed the literature involving *Sentinel Physician Networks*, using the Medline search strategy shown in Figure 1. Initially, 6691 articles matched the research criteria, with an increasing rate of publications over the past 20 years.

Most studies focused on clinical and organizational issues. Only a handful of studies (15) were related to EH, suggesting that only sporadic experience, based on some specific circumstances, has been acquired (Table 1). These findings show that knowledge and recognition of EH issues may not be a common practice amongst physicians, likely due to lack or training and/or awareness.

Nevertheless, some physicians have highlighted EH-related issues and shown how the environmental sensitivity of individual physicians and knowledge of their area, together with clinical expertise, can help detect critical situations and encourage appropriate actions to prevent recurrent threats (Table 2).

In summary, a real systematic initiative promoting clinical expertise in EH, including GCC and planetary health associated risks has not yet been developed. Hence, we believe it is time to act

PROPOSAL OF A FRAMEWORK OF A NETWORK OF SENTINEL PHYSICIANS

We are committed to developing a surveillance system comprised of sentinel FDs (including General Practitioners –GP-and Primary Care Paediatricians –PCPEDs-) in Italy, named *RIMSA* (*Rete Italiana Medici Sentinella per l'Ambiente- Italian Network of Sentinel Physicians for the Environment*), aimed at studying and monitoring the effects of environmental issues on population health, both to make a significant contribution towards raising awareness and informing the public, and to report any critical situations. It is well known that citizens strongly rely on FDs; thus there are good prospects that FDs, embedded within the social context where they practice, and within a professional network devoted to EH issues, can enhance their capacity to induce health promoting behaviour in the population.

• Furthermore, this proposal could:Promote a more



Table 1: List of publica	ntions and pre	esentations dealing v	vith Sentinel Practitioners and Environment.		
Author/s Year Country		Country	Topic		
BS. Levy ²⁰	1996	USA	Editorial: Holistic Approach to Surveillance		
I Devoux ²¹	2001	France	Wastewater reuse raises the question of health risk and the epidemiological surveys needed.		
J Litt et al. ²²	2004	USA	Survey of public health and environmental practitioners to uncover state and local health tracking needs and priorities		
A Hussa et al. ²³	2004	Switzerland	Estimates the scale of environmental medicine counselling in Switzerland by using two different data sources. The main source was the frequency of medical consultations due to environmental exposures in general practice the second using medical, psychological and environmental tools		
BS Schwartz et al. ²⁴	2005	USA	Editorial: community-based primary care providers must possess biomedical, epidemiologic, and environmental medicine skills,.		
A Flahault et al. ²⁵	2006	France	Description of <i>Réseau Sentinelles</i> : Database linkage with environmental information (e.g remote sensing, surface variables, environmental factors) will be facilitated, allowing for the evaluation of the role of climate change, or pollution involvement in disease.		
FH Johnston et al. ²⁶	2006	Australia	Investigates the relationship between particulate matter (PMPM10 and PM2.5) generated by vegetation fires and daily health outcomes in 251 adults and children with asthma over a 7-month period also recruited by GPs		
AJ Elliot et al. ²⁷	2006	UK	Investigates the association between impetigo, insect bites and air temperature: a retrospective 5-year study (1999-2003) using morbidity data collected from a sentinel general practice network database.		
SC Chen et al. ²⁸	2010	Taiwan	How to use a probability-based transmission modeling approach to examine the influen risk of infection virus in indoor environments. This was based on 10 years of data gathered from influenza-like illness sentinel physician and laboratory surveillance, and experimental viral shedding data in Taiwan.		
A Kolovos et al. ²⁹	2010	France	Development of model within an environmental health context which can be particularly important for prediction and decision-making in environmental health and risk studies, management, and planning, etc. Based on aggregated observations recorded by general physicians through the <i>Réseau Sentinelles</i>		
S. Medina ³⁰	2014	France	The analysis of drug consumption estimated an excess of about 5000 treatments by psychotropic drugs as a consequence of the explosion of a chemical plant (AZF) in Toulouse (2001)		
K Sebec et al ³¹	2014	USA	This experience allowed exploring the strengths and weaknesses of ambulatory Elettrinic Health Record (HER) data in post-disaster settings. Data from ambulatory EHR networks can augment existing surveillance streams by providing sentinel population snapshots on clinically available indicators in near real time.		
GL Nichols ³²	2014	Europe	They describe the surveillance systems (GP ones included), tracking tools, communication channels, information exchange and outputs in the light of environmenta and climatic drivers of infectious diseases		
C Klier et al. ³³	2016	Europe	This overview is aimed at providing current data on the incidence of Tick Borne Disease on longitudinal trends according to the environmental settings. Based on Sentinel physician results e.g. France, Netherlands, Switzerland. They indicate high reliability of collected data		
F Giroud et al ³⁴	2017	Madagascar	This study describes a system using various environmental and meteorological data with the support of new technologies to improve the performance of a sentinel Malaria-related data from 21 sentinel sites collected by Short Message Service which are automatically analysed to detect and forecast malaria trends and malaria outbreak alerts with automated feedback reports		

effective profile of FDs in Italy in the context of EH prevention. This is particularly relevant in Italy since it represents one of the areas of greatest impact due to GCC in Europe [44];

- Promote the integration of Departments of Primary Care and Prevention of the Local Health Authorities (LHA) with the Regional Agencies for the Environment (REPA);
- Re-affirm the role of FDs in promoting healthy and sustainable behaviours and lifestyles to cope with GCC health effects or any other kinds of EH threats.


We believe that our experience will be of value if shared


internationally with other Sentinel Practitioners Networks (see conclusions).


DESCRIPTION OF THE PROJECT CARRIED OUT IN ITALY

Goals achieved: a brief description of the results

Taking the opportunity offered by the project funded by the Italian Ministry of Health (MoH), within the preparatory process of the G7 health experts working group [45], a project aimed at promoting the definition and implementation of RIMSA has been developed.







An illustration of the overall project is provided in Figure 3.

A web site has been implemented [46], together with 3 courses in the south, centre and north of Italy (Taranto, Arezzo, Genoa), having trained 61 FDs from across the country.

An electronic survey to assess the trainees' learning needs

and wishes for future training initiatives has been carried out. A Moodle (acronym for *modular object-oriented dynamic*) platform for other training initiatives and an e-learning community of practice have been implemented [47].

A manual devoted to EH physicians willing to be involved in RIMSA is also being compiled.

It is noteworthy to point out that most of the professionals who attended the courses are now coordinating local actions, demonstrating the efficiency of the implementation of such a network. The final goal is to integrate these local experiences within a national and hopefully international context.

Sentinel Physician for the Environment (SPE): a professional profile.

- One critical issue has been properly investigated: the SPE must have a professional role and specific skills to fill the existing gap between health and environment at the community level. These skills can be summarised into two groups Clinical and epidemiological: To extrapolate significant signals of an environmentally-driven threat from patient reports; to select the most appropriate human biomarkers in the investigation of health problems, preferably choosing biomarkers of early subclinical effects; to know the main diseases related to local environmental threats.
- Advocacy towards prevention: to adopt the precautionary [48] and responsibility [49] principles dealing with EH threats; to influence and support the decision-making authorities; to inform the LHA about possible environmental problems that could represent risks for their patients' health; to collaborate with other disciplines/institutions; to educate his/her patients and their own family to promote an healthy lifestyle behaviour to cope with GCC and EH threats as a whole; to reduce the environmental impact of their medical activities as recommended by the green-health policy [50];

Information flows within the SPE network

A crucial effort will be devoted to defining the organized flow of information, to give a comprehensive answer to local environmental threats (e.g. incinerators, coke energy plants etc.) and/or global threats (e.g. GCC). As such, the following standard procedure has been conceived, according to a US-CDC guideline [51]:

- 1. Consulting the available bibliographic databases;
- Involvement of the RIMSA reference centre (RIMSA-RC) to check the "problem";
- Thereafter, the RIMSA-RC will ask for the collaboration of LHA and REPA;
- 4. If the "problem" is confirmed, a study will be designed with local RIMSA;
- 5. Finally, to reach conclusions on the presence of a local EH risk and make recommendations for action.

To achieve these goals without burdening the daily work of physicians, it will be necessary:



Table 2: Sentinel Physicians and Environment: some significant anecdotes.									
Name	Job	Year	Location	Outcomes	Due to	Alert consequences			
J. Snow ³⁵⁻³⁷	Anestesiologist	1854	London (UK)	V.Cholerae mortality	Drinkable water contaminated by sewage system	Change of tapping point upstream of London			
G.Franco ^{38,39}	Pediatrician	1980	Augusta-Priolo (SR) (I)	Birth malformations	industries leaking Mercury in the sea	Industry closure			
G. Porcile,	Oncologist	1980	Genoa (I)	Cancers	Incinerator	Incinerator closure			
MJ Gardner ⁴⁰	Professor Medical Science	1993	Sellafield (UK)	Limphoma	Nuclear Plant	Nuclear Plant Closure			
G. Costani ⁴¹	Family Doctor	1998	Mantova (I)	Soft tissue sarcoma	Industrial waste incinerator	Environmental Surveillance ongoing			
V. Cordiano ^{42,43}	Hematologist	2016	Treviso (I)	Higher mortality levels for some causes of death.	Teflon and Goretex production	Water safety limits set up			

- To set up a uniform data collection process, defining protocols, processes, standards, and core datasets, accounting for the ethical aspects of data collection, storage, consent and confidentiality, such as General Data Protection Regulation (GDPR) compliance;
- To implement a common EMR software, or at least procedures to allow comparable data extractions;
- To carry out geo-referencing (through GIS-based systems) of patients' addresses and accordingly of environmental data, which will be effectively managed (i.e., block chain; see below)

In line with Donabedian's principles [52], a continuous and systematic process of RIMSA Quality Assurance (QA) in terms of *structure, process, output and outcome indicators* will be established and implemented (Figure 3).

In conclusion, we emphasise that scientifically sound practice is key to supporting the strength of FDs' advocacy efforts.

General organizational criteria to develop and consolidate the network

The main challenge of the next phase of the *SPE* project will be to develop a framework to interact with other organisations.

One essential feature of RIMSA will be collaboration with other institutions and disciplines such as: LHA, REPA, Academia, municipalities and regional authorities, enterprises and trade unions, Inail (*Italian National Institute for Insurance against Accidents at Work*), the Italian Civil Protection and the media.

CONCLUSIONS

Lessons learned: RIMSA as a challenging exercise

Despite the definitions of the General Practice Network mentioned above (see page 8), whereby the aim of Sentinel networks is essentially of an epidemiological nature, the experience we are pursuing in Italy is quite challenging, because:

 The health effects of environmentally-driven phenomena are extremely complex in terms of how they can be

- interpreted (i.e. health risks and protective factors directly and indirectly linked to the environment);
- One essential facet of the General Practice Network is to couple the epidemiological mission with the opportunity to put in place the role of FDs in their community either at the individual or collective level, in the case of global threats (i.e. GCC).
- There are many organizations collecting and handling data not dealing with health (as was the case of the traditional Sentinel Practitioners network) but with environment [53,54], meteorology [55,56], occupation [57,58], etc.

What's next?

The ideas described here must now be implemented, which requires strong collaboration with institutional and professional bodies.

An active collaboration with the specialist training programmes (mainly in the field of General Practice and Paediatrics) must be pursued to develop the following didactic issues:

- What are the environmental-driven diseases? Which diseases could be identified at the local level by integrating epidemiological, toxicological, environmental knowledge?
- The concepts of hazard, risk and impact;
- Role of GPs and PCPEDs in promoting and protecting environmental health at an individual and collective level.

With reference to surveillance systems they may be enhanced through innovative approaches such as block chain technology [59-61].

Block chain could support the tracking of climate change and environmental health related health risks through electronic health records linked to other sources of data, such as indoor, outdoor and patient-wearable sensors, sensors that track the frequency, time and geo code of pharmacotherapy (e.g., Propeller Health: https://www.propellerhealth.com/), as well as patient-reported symptoms through smart phone or



smart watch applications directly linked to health records and disease-management software [62,63,64,65,66]. Furthermore, this initiative aims to be a forerunner also focusing on concerns that involve low-middle income countries. In particular, as it is essentially based on the socioeconomic, capacity and cultural/historical features in which the sentinel physicians operates, such an approach might be effectively implemented by coping it with other tragic conditions (i.e. malnutrition, waterborne diseases, infectious diseases, etc.), which should be taken into account not only for their direct health effects but also for others which indirectly influence the life and the health of people (ie migrations). Consequently, the WHO Department of Public Health and the Environmental and Social Determinants of Health declared its intention to collaborate in developing such an experience in an international perspective.

We would like to share our experience to raise interest in this field and to promote a "community" of experts and institutions to promote and support the essence of this concept.

Along this line, we involved some existing Sentinel Physician Networks operating across the world (in particular in Europe) such as in The Netherland, the UK, Germany and France (Brussels June 27th, 2018). Collectively, we will develop an international clearinghouse for FDs, public health practitioners and researchers on how to advance and enhance national SPEs' capacity. In other words we aim to create an international network to support this network.

In conclusion, it is noteworthy to recall what Van Royen et al., stated in 2010: "further research in the area of personcentred care, comprehensive and holistic approach, should focus on: understanding how social, cultural and environmental circumstances that may have an effect on different aspects of health." [67].

FUNDING DETAILS

This paper was supported by the Agenzia Italiana per la Cooperazione e lo Sviluppo (AICS) within the project "Educare alla cittadinanza e alla salute globale".

COLLABORATION

B.Aimone, C.Alicino, V. Andrei, A. Autunno, M. Baccini, F.Balestreri ,L.Barberio, F.Bartolini, T.Battista, G. Bianchi, N.Bianco, D.Bottero, S.Borgo, C. Bregant, M.Calgaro, S.Catalano, F. Cavasin, M. Ciasullo, T. Cimolino, L. Cori, A. Cuffari, S. Dalmonte, M.De Vita, G. Del Gaiso, F. Di Bello, A. Di Benedetto, A. Faggioli, I. Ferrari, A.Fortunato, E.Franceschini, M.Godone, M.Greco, F. Laghi, M.Lepido, F.Licordari, L. Lucentini , F.Macchia, P. Maddalena, S. Mangia, G. Miserotti, P. Milo, S. Miozzo G.Montalbano, G.Occhini, A.Panajia, S.Parisi, S. Patania, R.Pedale, P.Pedote, G.Piana, S.Piussi, G.Pulito, M.Restuccia, R.Sammarco, R.Seller, L.Terzano, E.Trastulli, B.Travella, P.L. Tucci M.Turolla, A. Valente, G.Vallefuoco, F.Viganò, G.Zagami, R.P.Zambelli

ACKNOWLEDGEMENT

This publication has been supported by the "Agenzia Italiana per la cooperazione e lo sviluppo".

The authors' special thanks go to CUAMM for its precious

collaboration

"This publication has been made possible with the contribution of the Italian Agency for Development and Cooperation. The authors are the only persons responsible for the contents of this publication which do not necessarily represent the Agency's point of view"

REFERENCES

- Prüss-Ustün A, Wolf J, Corvalán C, Bos R, Neira M. Preventing disease through healthy environments. A global assessment of the burden of disease from environmental risks. Geneva, Switzerland: WHO. 2016.
- Nick Watts, Markus Amann, Sonja Ayeb-Karlsson. The Lancet 1-50.
 2017.
- Hales S, Kovats S, Lloyd S, Campbell-Lendrum D. Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization. 2014.
- Hamaoui-Laguel. Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe. Nat Clim Change. 2015; 5: 766–771.
- BQ Li , Wang J, Lavigne E, Gasparrini A, Copes R, Yagouti A, et al. Hospitalizations from Hypertensive Diseases, Diabetes, and Arrhythmia in Relation to Low and High Temperatures: Population-Based Study. Scientific Reports. 2016; 6: 30283.
- 6. Hopp S, Dominici F, Bobb JF. Medical diagnoses of heat wave-related hospital admissions in older adults. Prev Med. 2018; 110: 81-85.
- 7. Turner JB. The effects of indoor and outdoor temperature on metabolic rate and adipose tissue the Mississippi perspective on the obesity epidemic. Rev Endocr Metab Disord. 2016; 17: 61-71.
- 8. Hajat S, Haines A, Sarran C, Sharma A, Bates C, Fleming LE. The effect of ambient temperature on type2-diabetes: case-crossover analysis of 4+ million GP consultations across England Environmental Health. 2017; 16: 73.
- 9. Dain K, Handley L. Diabetes and climate change--two interconnected global challenges. Diabetes Res Clin Pract. 2012; 97: 337-339.
- 10. Bunker A, Wildenhain J, Vandenbergh A, Henschke N, Rocklöv J, Hajat S, et al. Effects of air temperature on climate-sensitive mortality and morbidity outcomes in the elderly; a systematic review and meta-analysis of epidemiological evidence. EBioMedicine. 2016; 6: 258-268.
- 11. Cho J, Choi YJ, Sohn J, Suh M, Cho SK, Ha KH, et al. Ambient ozone concentration and emergency department visits for panic attacks. J Psychiatr Res. 2015; 62: 130-135.
- 12. Kirrane EF, Bowman C, Davis JA, Hoppin JA, Blair A, Chen H, et al. Associations of Ozone and PM2.5 Concentrations With Parkinson's Disease Among Participants in the Agricultural Health Study. J Occup Environ Med. 2015; 57: 509-517.
- 13. Aström C, Rocklöv J, Hales S, Béguin A, Louis V, Sauerborn R. Potential distribution of dengue fever under scenarios of climate change and economic development. Ecohealth. 2012; 9: 448-454.
- 14. Schaffner F, Mathis A. Dengue and dengue vectors in the WHO European region: past, present, and scenarios for the future. Lancet Infect Dis. 2014; 14: 1271-1280.
- 15. Hummers-Pradier E, Beyer M, Chevallier P, Eilat-Tsanani S, Lionis C, Peremans L. The Research Agenda for General Practice/Family Medicine and Primary Health Care in Europe. Part 1. Background and methodology. 2009; 15: 243–250.
- 16. Malkhazova SM, Mironova VA, Kotova TV, Shartova NV, Orlov DS. Natural-focal diseases: mapping experience in Russia Int J Health



- Geogr. 2014; 13: 21.
- 17. Busani L, Platonov AE, Ergonul O, Rezza G. How to Tackle Natural Focal Infections: From Risk Assessment to Vaccination Strategies. Adv Exp Med Biol. 2017; 972: 7-16.
- 18.Xie E, Falceto de Barros E, Abelsohn A, Tetelbom Stein A, Haines A. Challenges and opportunities in planetary health for primary care providers. 2018; 2.
- 19. Van Casteren V. Eurosentinel. In: Fracchia GN, Theofilatou M (eds) Health services research. Commission of the European Communities, Directorate-General for Science, Research and Development. IOS Press, Amsterdam.1993; 253–262.
- 20. Levy BS. Toward a Holistic Approach to Public Health Surveillance. Am J Public Health. 1996; 86: 624-625.
- 21. Devaux I, Gerbaud L, Planchon C, Bontoux J, Glanddier PY. Infectious risk associated with wastewater reuse: an epidemiological approach applied to the case of Clermont-Ferrand, France. Water Sci Technol. 2001: 43: 53-60.
- 22. Litt J, Tran N, Chossek Malecki K, Neff R, Resnick B, Burke T. Identifying Priority Health Conditions, Environmental Data, and Infrastructure Needs: A Synopsis of the Pew Environmental Health Tracking Project. Environ Health Perspect. 2004; 112: 1414–1418.
- 23. Hussa A, Küchenhoffb J, Bircherc A, Niedererd M, Trempe J, Waeberf R, Braun-Fahrländera C, for the Swiss Sentinel Surveillance Networkg. Are environmental medicine problems relevant in Switzerland? SWISS MED WKLY. 2004; 134: 500–507.
- 24. Schwartz BS, Rischitelli G, Hu H. The Future of Environmental Medicine in Environmental Health Perspectives: Where Should We Be Headed? Environmental Health Perspectives. 2005; 113: 574-576.
- 25. Flahault A, Blanchon T, Dorléans Y, Toubiana L, Vibert JF, Valleron AJ. Virtual surveillance of communicable diseases: a 20-year experience in France Statistical Methods in Medical Research. 2006; 15: 413–421.
- 26. Johnston FH, Webby RJ, Pilotto LS, Bailie RS, Parry DL, Halpin SJ. Vegetation fires, particulate air pollution and asthma: A panel study in the Australian monsoon tropics. Int J Environ Health Res. 2006; 16: 391–404.
- 27. Elliot AJ. Cross KW, Smith GE, Burgess IF, Fleming DM. The association between impetigo, insect bites and air temperature: a retrospective 5-year study (1999-2003) using morbidity data collected from a sentinel general practice network database. Family Practice. 2006; 23: 490-496.
- 28. Chen SC, Liao CM. Probabilistic indoor transmission modelling for influenza (sub)type viruses. The J Infect. 2010; 60: 26-35.
- 29. Kolovos A, Angulo JM, Modis K, Papantonopoulos G, Wang JF, Christakos G. Model-driven development of covariances for spatiotemporal environmental health assessment. Environ Monit Assess. 2013; 185: 815-831.
- 30. Medina S. The experience of developing environmental-health dimensions of routine surveillance data systems Environmental Public Health Tracking: Practical Methods for Priority Setting and Evaluation. ISEE pre-conference workshop. 2013.
- 31. Sebek K, Jacobson L, Wang J, Newton-Dame R, Singer J. Assessing Capacity and Disease Burden in a Virtual Network of New York City Primary Care Providers Following Hurricane Sandy J Urban Health. 2014: 91: 615-622.
- 32. Nichols GL, Andersson Y, Lindgren E, Devaux I, Semenza JC. European Monitoring Systems and Data for Assessing Environmental and Climate Impacts on Human Infectious Diseases Int. J. Environ. Res. Public Health. 2014; 11: 3894-3936.

- 33. Klier C, S. O'Connell, Kleinemeier C, Smith R. Surveillance of Lyme Borreliosis in Europe: Trend analysis 1999 through 2014 Health—exploring complexity: an interdisciplinary systems approach HEC2016 28 August–2 September 2016, Munich, Germany.
- 34. Girond F, Randrianasolo L, Randriamampionona L, Rakotomanana F, Randrianarivelojosia M, Ratsitorahina M. Analysing trends and forecasting malaria epidemics in Madagascar using a sentinel surveillance network: a web-based application Malar J. 2017; 16: 72.
- 35. John S. On the Mode of Communication of Cholera. London: John Churchill. 1849.
- 36.Paneth N, Fine P. The art of medicine The singular science of John Snow. 2013; 381.
- 37.AM Lilienfeld, DE Lilienfeld, Fondamenti di Epidemiologia, pag 36 Ed Piccin 1986.
- 38. Bianchi F, Bianca S, Linzalone N, Madeddu A. Surveillance of congenital malformations in Italy: an investigation in the province of Siracusa Epidemiol Prev. 2004; 28: 87-93.
- 39.La Syndial offre «somme di ristoro» alle donne di Augusta-Priolo che hanno abortito o partorito figli con gravi malformazioni Epidemiol Prev. 2006; 30: 76-77.
- 40. Gardner MJ. Father's occupational exposure to radiation and the raised level of childhood leukemia near the Sellafield nuclear plant. Environ Health Perspect. 1991; 94: 5–7.
- 41. Costani G. Incidenza anomala di sarcomi dei tessuti molli a Mantova. Epidemiol Prev. 1998; 22: 1.
- 42. Cordiano V, Storti M, Bai E, Crosignani P. Perfluoroalkyl substances in groundwater in Veneto Region (Northern Italy): a new Seveso case? Epidemiol Prev. 2017; 41: 148-148.
- 43. Mastrantonio M, Bai E, Uccelli R, Cordiano V, Screpanti A, Crosignani P. Drinking water contamination from perfluoroalkyl substances (PFAS): an ecological mortality study in the Veneto Region, Italy. Eur J Public Health. 2017; 28: 1–6.
- 44. Climate and Health Country Profile, Italy. World Health Organisation, Geneva. 2018.
- 45.http://www.g7italy.it/sites/default/files/documents/FINAL_G7_Health_Communiqu%C3%A8_Milan_2017_0.pdf
- 46. http://www.isde.it/rete-italiana-medici-sentinella-lambiente/
- 47. http://research.moodle.net/mod/data/view.php?d=1&rid=154
- 48.http://www.un.org/documents/ga/conf151/aconf15126-1annex1.htm
- 49. Hans Jonas Das Prinzip Verantwortung: Versuch einer Ethik für die technologische Zivilisation. Frankfurt/M., 1979. New edition: Suhrkamp Taschenbuch, 1984.
- 50.https://noharm.org/
- 51.Beth A, Anderson H, Blackmore C, Bove FJ, Condon SK, Eheman CR, et al. "Investigating Suspected Cancer Clusters and Responding to Community Concerns: Guidelines from CDC and the Council of State and Territorial Epidemiologists." Morbidity and Mortality Weekly Report: Recommendations and Reports. JSTOR. 2013; 62: 1–24.
- 52. Donabedian A. Quality of Care, How Can It Be Assessed? JAMA. 1988; 260: 1743-1748.
- 53.https://www.eea.europa.eu/
- 54.http://www.isprambiente.gov.it/it
- 55.https://www.ecmwf.int/



- 56.https://www.cmcc.it/it/
- 57. http://www.ilo.org/global/lang--en/index.htm
- 58. https://www.inail.it/cs/internet/home.html
- 59. Iansiti M, Lakhana KR. The Truth About Blockchain. Harvard Business Review. 2017.
- 60. Nakamoto S. 2998. Bitcoin: A peer-to-peer electronic cash system.
- 61. Hashed Health. Blockchain Public Health Surveillance.
- 62.Stagnaro C, Freed Associates. White Paper: Innovative Blockchain Uses in Health Care.
- 63.Brodersen C, Kalis B, Leong C, Mitchell E, Trusscott A, Accenture LLP. Blockchain: Securing a New Health Interoperability Experience.

Accenture, 2016.

- 64. Thomas H. A case study in blockchain health care innovation. Int J Curr Res 2017; 9: 60587-60588.
- 65. P Lauriola, G Leonardi, E Righi, T Bayleyegn, A Schnall, J Malilay, S Louca, B Behbod, Natural Disaster Environmental Health Preparedness In: Encyclopedia of Environmental Health, Elsevirer Ed. 2018.
- 66.Kyriakoudes G, Soulla L, Behbod B. Cyprus's new National Health Service and future European health. Lancet. 2018.
- 67. van Royen P, Beyer M, Chevallier P, Eilat-Tsanani S, Lionis C, Peremans L. The research agenda for general practice/family medicine and primary health care in Europe. Part 5: Needs and implications for future research and policy. Eur J Gen Pract. 2010; 16: 244–248.

Cite this article

Lauriola P, Pegoraro S, Serafini A, Murgia V, Di Ciaula A, et al. (2018) The Role of General Practices for Monitoring and Protecting the Environment and Health. Results and Proposals of the Italian Project Aimed at Creating an "Italian Network of Sentinel Physicians for the Environment" (Rimsa) within an International Perspective. J Family Med Community Health 5(5): 1160.